

## Quick Reference to Transfusion Reaction Action

For ALL suspected blood component/product transfusion reactions

**STOP** the transfusion. Do NOT infuse the remaining blood component/product in the line.

- |  |  |
|--|--|
| <ol style="list-style-type: none"> <li>1. Do not discard product.</li> <li>2. Disconnect the set from the IV site, while maintaining sterility of the blood product and IV set.             <ul style="list-style-type: none"> <li>• Using new IV set, administer Normal Saline (or D5W for IVIG infusion) at TKVO.</li> </ul> </li> <li>3. Contact MD/designate for medical assessment/treatment</li> <li>4. Assess the patient's vital signs at least every 15 minutes until patient is stable</li> <li>5. Perform visual assessment of product</li> </ol> | <ol style="list-style-type: none"> <li>6. Check for clerical discrepancies             <ul style="list-style-type: none"> <li>• Confirm patient identification</li> <li>• Confirm patient demographics and all documentation matches:                 <ul style="list-style-type: none"> <li>○ ID band/health card</li> <li>○ Physician order in chart</li> <li>○ Tag on product</li> <li>○ Label on product</li> <li>○ Transfusion Medicine Results Report</li> </ul> </li> </ul> </li> <li>7. Notify blood bank/lab</li> </ol> |
|--|--|

Table 1: Clinical Signs and Symptoms (S/S)	Send to Blood Bank			"Ongoing Transfusion Care"
	Transfusion Reaction Form	Sealed Blood Product	Patient Samples	
IVIG related, mild transient S/S- side effects that resolve with reduced flow rate or medication	No	No	None	May restart the transfusion at a slower rate with appropriate medication and frequent vital signs <b>if</b> ordered by the physician after consultation regarding the patient's condition  Direct observation for 15 minutes following transfusion re-establishment
Urticaria or pruritus with any blood component/product	Yes	No	None	
Oral temperature rise greater than 1°C <b>AND</b> Temperature less than 39°C <b>AND</b> onset greater than 10 minutes into transfusion	Yes	No	None	
IVIG related S/S that are moderate or severe or unresponsive to <b>"Ongoing Transfusion Care"</b>	Yes	Yes	None	<b>DO NOT restart transfusion</b>
Suspected bacterial contamination (see Table 2 for S/S)	Yes	Yes (avoid contamination of product)	2 EDTA (purple tubes) <b>AND</b> Patient Blood Cultures are recommended	
All other unexpected signs or symptoms with any blood component/product	Yes	Yes (avoid contamination of product)	2 EDTA (purple tubes)	

<b>Table 2:</b> <b>SUSPECT</b> Bacterial Contamination of the blood component/product <b>IF</b> patient has <b>Signs or Symptoms A,B,C or D</b> as follows:		
<b>A.</b> Temperature rise more than 1 °C <b>AND</b> greater than 39°C	<b>B.</b> Temperature rise greater than 1°C <b>AND</b> between 38°C <b>AND</b> 39°C with: *Rigors   *Hypotension   *Tachycardia   *Dyspnea and/or *Shock	<b>C.</b> Temperature rise not responding to antipyretic <b>AND/OR</b>  <b>D.</b> Suspicion of sepsis in absence of fever

**NOTE: Consult Transfusion Medicine MD on Call if replacement blood products are required urgently.**